

A NEW HIGH-REDSHIFT LYMAN α EMITTER: POSSIBLE SUPERWIND GALAXY AT $Z = 5.69$ ^{1, 2}

MASARU AJIKI ³, YOSHIAKI TANIGUCHI ³, TAKASHI MURAYAMA ³, TOHRU NAGAO ³, SYLVAIN
 VEILLEUX ⁴, YASUHIRO SHIOYA ³, SHINOBU S. FUJITA ³, YUKO KAKAZU ⁵, YUTAKA KOMIYAMA ⁶,
 SADANORI OKAMURA ^{7,8}, DAVID B. SANDERS ⁵, SHINKI OYABU ⁹, KIMIAKI KAWARA ⁹, YUICHI
 OHYAMA ⁶, MASANORI IYE ¹⁰, NOBUNARI KASHIKAWA ¹⁰, MICHITOSHI YOSHIDA ¹¹, TOSHIYUKI
 SASAKI ⁶, GEORGE KOSUGI ⁶, KENTARO AOKI ⁹, TADAFUMI TAKATA ⁶, YOSHIHIKO SAITO ¹⁰, KOJI
 S. KAWABATA ¹⁰, KAZUHIRO SEKIGUCHI ⁶, KIICHI OKITA ⁶, YASUHIRO SHIMIZU ¹¹, MOTOKO INATA
⁹, NOBORU EBIZUKA ¹², TOMOHIKO OZAWA ¹³, YASUSHI YADOUMARU ¹³, HIROKO TAGUCHI ¹⁴,
 HIROYASU ANDO ⁶, TETSUO NISHIMURA ⁶, MASAHIKO HAYASHI ⁶, RYUSUKE OGASAWARA ⁶, &
 SHIN-ICHI ICHIKAWA ¹⁰

Accepted for publication in the Astrophysical Journal

ABSTRACT

During the course of our deep optical imaging survey for Ly α emitters at $z \approx 5.7$ in the field around the $z = 5.74$ quasar SDSSp J104433.04-012502.2, we have found a candidate strong emission-line source. Follow-up optical spectroscopy shows that the emission line profile of this object is asymmetric, showing excess red-wing emission. These properties are consistent with an identification of Ly α emission at a redshift of $z = 5.687 \pm 0.002$. The observed broad line width, $\Delta v_{\text{FWHM}} \simeq 340 \text{ km s}^{-1}$ and excess red-wing emission also suggest that this object hosts a galactic superwind.

Subject headings: galaxies: individual (LAE J1044–0130) — galaxies: starburst — galaxies: formation

1. INTRODUCTION

Recent progress in deep optical imaging with 8-10 m class telescopes has enabled new searches for star-forming galaxies beyond redshift 5. In particular, imaging surveys using narrow-passband filters have proved to be a particularly efficient way to find such galaxies (Hu & McMahon 1996; Cowie & Hu 1998; Steidel et al. 2000; Kudritzki et al. 2000; Hu et al. 2002). Indeed the most distant Ly α emitter known to date is HCM 6A at $z = 6.56$ (Hu et al. 2002), and more than a dozen Ly α emitters beyond $z = 5$ have been discovered (Dey et al. 1998; Spinrad et al. 1998; Weymann et al. 1998; Hu et al. 1998, 1999, 2002; Dawson et al. 2001, 2002; Ellis et al. 2001), most by using this technique.

One interesting object is J123649.2+621539 at $z = 5.190$ which was found serendipitously in the HDF-North flanking fields (Dawson et al. 2002). Its Ly α emission-line profile shows a sharp blue cutoff and broad red wing emission, both of which are often observed in star-forming systems with prominent wind outflows. These features are also ex-

pected from radiative transfer in an expanding envelope. Therefore, Dawson et al. (2002) suggested that the Ly α profile of J123649.2+621539 is consistent with a superwind with a velocity of $\sim 300 \text{ km s}^{-1}$. Galactic superwinds are now considered to be one of the key issues for understanding the interaction and evolution of both galaxies and intergalactic matter (e.g. Heckman 1999; Taniguchi & Shioya 2000). In order to improve our knowledge of galactic superwinds at high redshift, a large sample of superwind candidates at $z > 3$ is needed. During the course of our new search for Ly α emitters at $z \approx 5.7$, we have found a candidate superwind galaxy at $z = 5.69$. In this *Letter*, we report its observed properties. We adopt a flat universe with $\Omega_{\text{matter}} = 0.3$, $\Omega_{\Lambda} = 0.7$, and $h = 0.7$ where $h = H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1})$ throughout this *Letter*.

2. OPTICAL DEEP IMAGING

We have carried out a very deep optical imaging survey for faint Ly α emitters in the field surrounding the quasar SDSSp J104433.04–012502.2 at redshift 5.74¹⁵ (Fan et al.

¹Based on data collected at the Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

²Based on data collected at the W. M. Keck Observatory

³Astronomical Institute, Graduate School of Science, Tohoku University, Aramaki, Aoba, Sendai 980-8578, Japan

⁴Department of Astronomy, University of Maryland, College Park, MD 20742, USA

⁵Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

⁶Subaru Telescope, National Astronomical Observatory, 650 N. A'ohoku Place, Hilo, HI 96720, USA

⁷Department of Astronomy, Graduate School of Science, University of Tokyo, Tokyo 113-0033, Japan

⁸Research Center for the Early Universe, School of Science, University of Tokyo, Tokyo 113-0033, Japan

⁹Institute of Astronomy, Graduate School of Science, University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan

¹⁰National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

¹¹Okayama Astrophysical Observatory, National Astronomical Observatory, Kamogata-cho, Asakuchi-gun, Okayama 719-0232, Japan

¹²Communications Research Laboratory, 4-2-1 Nukui-Kitamachi, Koganei, Tokyo 184-8795, Japan

¹³Misato Observatory, 180 Matsugamine, Misato-cho, Amakusa-gun, Wakayama, 640-1366, Japan

¹⁴Department of Astronomy and Earth Sciences, Tokyo Gakugei University, 4-1-1 Nukui-Kitamachi, Koganei, Tokyo 184-8501, Japan

¹⁵The discovery redshift was $z = 5.8$ (Fan et al. 2000). Since, however, the subsequent optical spectroscopic observations suggested a bit lower redshift; $z = 5.73$ (Djorgovski et al. 2001) and $z = 5.745$ (Goodrich et al. 2001), we adopt $z = 5.74$ in this *Letter*.

2000; Djorgovski et al. 2001; Goodrich et al. 2001), using the prime-focus wide-field camera, Suprime-Cam (Miyazaki et al. 1998) on the 8.2 m Subaru Telescope (Kaifu 1998) on Mauna Kea. Suprime-Cam consists of ten $2k \times 4k$ CCD chips and provides a very wide field of view $-34' \times 27'$ ($0.2 \text{ arcsec pixel}^{-1}$). In this survey, we used the narrow-passband filter, NB816, centered on $8160 \text{ \AA}$ with a passband of $\Delta\lambda_{\text{FWHM}} = 120 \text{ \AA}$; the central wavelength corresponds to a redshift of 5.72 for $\text{Ly}\alpha$ emission. We also used broad-passband filters, B , R_C , I_C , and z' . A summary of the imaging observations is given in Table 1. All observations were done under photometric conditions, and the seeing was between $0''.7$ and $1''.3$ arcsec during the entire run. Photometric and spectrophotometric standard stars used in the flux calibration are SA101 for the B , R_C , and I_C data, and GD 50, GD 108 (Oke 1990), and PG 1034+001 (Massey et al. 1996) for the NB816 data. The z' data were calibrated by using the magnitude of SDSSp J104433.04–012502.2 (Fan et al. 2000); since any quasar is a potentially variable object, the photometric calibration of the z' data may be more unreliable than those of the other-band data.

To avoid delays in obtaining follow-up spectroscopy, we analyzed only two of the CCD chips, one of which included the quasar SDSSp J104433.04–012502.2. The individual CCD data were reduced and combined using IRAF and the mosaic-CCD data reduction software developed by Yagi et al. (2002). The total size of the reduced subfield is $11'.67 \times 11'.67$, corresponding to a total solid angle of $\approx 136 \text{ arcmin}^2$. The volume probed by the NB816 imaging has (co-moving) transverse dimensions of $27.56 h_{0.7}^{-1} \text{ Mpc} \times 27.56 h_{0.7}^{-1} \text{ Mpc}$, and the FWHM half-power points of the filter correspond to a co-moving depth along the line of sight of $44.34 h_{0.7}^{-1} \text{ Mpc}$ ($z_{\text{min}} \approx 5.663$ and $z_{\text{max}} \approx 5.762$; note that the transmission curve of our NB816 filter has a Gaussian-like shape). Therefore, a total volume of $33,700 h_{0.7}^{-3} \text{ Mpc}^3$ is probed in our NB816 image.

Source detection and photometry were performed using SExtractor version 2.2.1 (Bertin & Arnouts 1996). Our detection limit (a 3σ detection with a $2''$ diameter aperture) for each band is summarized in Table 1. For source detection in the NB816 image, we used a criterion that a source must be an object with at least a 13-pixel connection at the 5σ level. [Note: Given the pixel resolution of $0.2 \text{ arcsec pixel}^{-1}$, an extended source observed in $\sim 1 \text{ arcsec}$ seeing must be observed as a source with more than a 13-pixel connection.] Adopting the criterion for the NB816 excess, $I_C - \text{NB816} > 1.0 \text{ mag}$, we have found a strong emission-line source located at $\alpha(\text{J2000.0})=10^{\text{h}} 44^{\text{m}} 32^{\text{s}}$ and $\delta(\text{J2000.0})=-01^\circ 30' 34''$ (hereafter LAE J1044–0130). In this *Letter* we report on this source.

The sky position of LAE J1044–0130 is shown in Fig. 1. The optical thumb-nail images of LAE J1044–0130 are given in Fig. 2. As shown in this figure, LAE J1044–0130 is clearly seen only in the NB816 image; the observed equivalent width is $EW_{\text{obs}} > 310 \text{ \AA}$. The NB816 image reveals that LAE J1044–0130 is spatially extended. It is interesting to note that this object shows not a circular shape but an irregular shape. The angular diameter is 2.3 arcsec (above the 2σ noise level). The size of the point spread function in the NB816 image is 0.90 arcsec . Correcting for this spread, we obtain an angular diameter of

2.1 arcsec for the object. In the cosmology adopted here, this corresponds to a diameter of $d \simeq 12.4 h_{0.7}^{-1} \text{ kpc}$.

3. OPTICAL SPECTROSCOPY

In order to investigate the nature of LAE J1044–0130, we obtained optical spectroscopy. First, we used the Subaru Faint Object Camera And Spectrograph (FOCAS; Kashikawa et al. 2000) with the lowest spectral resolution grism, 150 lines mm^{-1} , blazed at $\lambda=6500 \text{ \AA}$ together with an order-cut filter SY47 on 2002, March 11 (UT). The wavelength coverage was $\sim 4900 \text{ \AA}$ to $9400 \text{ \AA}$. The use of an 0.8 arcsec -wide slit gave a spectroscopic resolution of $R \sim 400$ at $8000 \text{ \AA}$. The total integration time was 1800 seconds. The spectrum is shown in the upper panel of Fig. 3. We detected a single emission line at $\lambda \approx 8130 \text{ \AA}$. This observation was done under photometric conditions, and we used the spectroscopic standard star, Feige 34 (Massey et al. 1988) to calibrate the spectrum. The emission line flux was calculated to be $(1.5 \pm 0.3) \times 10^{-17} \text{ ergs s}^{-1} \text{ cm}^{-2}$. A second spectrum with FOCAS was obtained on 2002, March 13 (UT) in non-photometric conditions. This time, we used the 300 lines mm^{-1} grating blazed at $7500 \text{ \AA}$ together with the same order cut filter SY47, giving a spectral resolution $R \simeq 1000$ with the same 0.8 arcsec -wide slit. This setting gave the same wavelength coverage as the lower resolution observation. Again the total integration time was 1800 seconds. The spectrum is shown in the lower panel of Fig. 3. The emission-line profile marginally shows the sharp cutoff at wavelengths shortward of the line peak. However, the spectral resolution is insufficient to confirm this feature.

To clarify the shape of the line profile seen in the FOCAS observations, additional observations were obtained with Keck/Echelle Spectrograph and Imager (ESI; Sheinis et al. 2000) in echellette mode using a $1''$ slit, which provided a spectral resolution of $R \simeq 3400$ at $\lambda=8000 \text{ \AA}$. Two 1800-second integrations were obtained in photometric conditions on 2002, March 15 (UT). The spectra were calibrated using the spectroscopic standard stars Feige 34 and HZ 44 (Massey et al. 1988). The combined spectrum is shown in the middle panel of Fig. 4. Again only one emission line, at $\lambda \approx 8130 \text{ \AA}$ is found in the ESI optical spectrum, which covers the wavelength range $4000 \text{ \AA}$ to $9500 \text{ \AA}$. The emission-line profile at $8130 \text{ \AA}$ is confirmed to slightly asymmetric, showing a cutoff at wavelengths shortward of the line peak.

Although a single emission line in the wavelength range $4000 \text{ \AA}$ to $9500 \text{ \AA}$ may alternatively be identified as $[\text{O III}]\lambda 3727$, the fact that the observed profile appears to be skewed suggests that this line is in fact $\text{Ly}\alpha$ emission (e.g. Stern et al. 2000; Dawson et al. 2002). Therefore, we conclude that the emission line at $8129.3 \pm 3.0 \text{ \AA}$ must be $\text{Ly}\alpha$, which then yields a redshift of 5.687 ± 0.002 . The rest-frame equivalent width of the putative $\text{Ly}\alpha$ emission is estimated to be $EW_0 > 46 \text{ \AA}$. The observed $\text{Ly}\alpha$ flux is $(1.49 \pm 0.33) \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$. This appears consistent with our FOCAS observations made on 2002, March 11.

4. RESULTS AND DISCUSSION

4.1. Star Formation and Superwind Activities

The observed Ly α flux is $f(\text{Ly}\alpha) = (1.49 \pm 0.33) \times 10^{-17}$ ergs cm $^{-2}$ s $^{-1}$ based on the ESI spectrum. Given the cosmology adopted in this *Letter*, we obtain an absolute Ly α luminosity of $L(\text{Ly}\alpha) \simeq (5.3 \pm 1.2) \times 10^{42} h_{0.7}^{-2}$ ergs s $^{-1}$. This Ly α luminosity is comparable to those of other $z > 5$ galaxies; 1) 3.3×10^{42} ergs s $^{-1}$ for HCM 6A at $z = 6.56$ (Hu et al. 2002), 2) 6.1×10^{42} ergs s $^{-1}$ for SSA22-HCM1 at $z = 5.74$ (Hu et al. 1999), 3) 3.4×10^{42} ergs s $^{-1}$ for HDF 4-473.0 at $z = 5.60$ (Weymann et al. 1998), and 4) 8.5×10^{42} ergs s $^{-1}$ for J123649.2+621539.5 at $z = 5.19$ (Dawson et al. 2002). Note that all the above luminosities are estimated by using the same cosmology as that used here.

We note that approximately half of the intrinsic Ly α emission from LAE J1044-0130 could be absorbed by intergalactic atomic hydrogen (e.g., Dawson et al. 2002). In order to reproduce the observed Ly α emission-line profile a two-component profile fit was made using the following assumptions: 1) the intrinsic Ly α emission line profile is Gaussian, and 2) the optical depth of the Ly α absorption increases with decreasing wavelength shortward of the rest-frame Ly α peak. The resulting fit is shown in Figure 4 (thick curve), which corresponds to the following emission and absorption line parameters – 1) Ly α emission: the line center, $\lambda_{c,\text{em}} = 8030.70$ Å, the line flux, $f_{\text{em}} \simeq 2.42 \times 10^{-17}$ ergs s $^{-1}$ cm $^{-2}$, and the line width, $FWHM_{\text{em}} \simeq 650$ km s $^{-1}$; 2) Ly α absorption: the line center, $\lambda_{c,\text{abs}} = 8122.73$ Å, the optical depth at the absorption center, $\tau_{\text{abs}} \simeq 9.85$, and the line width, $FWHM_{\text{abs}} \simeq 175$ km s $^{-1}$. This analysis suggests that the total Ly α emission-line flux amounts to 1.73×10^{-17} ergs s $^{-1}$ cm $^{-2}$. This is larger by a factor of 1.16 than the observed flux, giving a total Ly α luminosity of $L(\text{Ly}\alpha) \sim 6.1 \times 10^{42} h_{0.7}^{-2}$ ergs s $^{-1}$. We then estimate the star formation rate of LAE J1044-0130. Using the relation $SFR = 9.1 \times 10^{-43} L(\text{Ly}\alpha) M_{\odot} \text{yr}^{-1}$ (Kennicutt 1998; Brocklehurst 1971) and the total Ly α luminosity, we obtain $SFR = 5.6 \pm 1.1 h_{0.7}^{-2} M_{\odot} \text{yr}^{-1}$. Note that SFR

may be overestimated because part of the Ly α emission may arise from shock-heated gas if the superwind interpretation is applicable to this object.

Although the signal-to-noise ratio of our ESI spectrum is not high enough to analyze the profile shape in great detail, the presence of the excess red-wing emission seems secure (Fig. 4). The FWHM of the Ly α emission is measured to be 340 ± 110 km s $^{-1}$ and the full width at zero intensity (FWZI) is estimated to be 890 ± 110 km s $^{-1}$. These properties are similar to those of the Ly α emitter at $z = 5.190$, J123649.2+621539, found by Dawson et al. (2002).

4.2. Comments on Possible Association with the Quasar SDSSp J104433.04-012502.2 and the Lyman Limit System at $z = 5.72$

The observed redshift of LAE J1044-0130, $z = 5.687$, is close both to the quasar redshift, $z = 5.74$ (see footnote 15) and to that of a Lyman limit system (LLS) at $z_{\text{LLS}} = 5.72$ in the quasar spectrum reported by Fan et al. (2000). The redshift difference between LAE J1044-0130 and the quasar corresponds to the velocity difference of $\Delta v \approx 2370$ km s $^{-1}$ and that between LAE J1044-0130 and the LLS corresponds to $\Delta v \approx 1476$ km s $^{-1}$. The angular distance between LAE J1044-0130 and the quasar is approximately 330 arcsec, giving a co-moving separation of $\sim 13 h_{0.7}^{-1}$ Mpc. This separation seems too large to identify LAE J1044-0130 as a counterpart of the LLS. It seems also unlikely that LAE J1044-0130 is associated with the large-scale structure in which the quasar SDSSp J104433.04-012502.2 resides.

We would like to thank both the Subaru and Keck Telescope staff for their invaluable help. We would also like to thank the referee for useful comments. This work was financially supported in part by the Ministry of Education, Culture, Sports, Science, and Technology (Nos. 10044052, and 10304013).

REFERENCES

- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393
 Brocklehurst, M. 1971, MNRAS, 153, 471
 Cowie, L. L., & Hu, E. M. 1998, AJ, 115, 1319
 Dawson, S., Spinrad, H., Stern, D., Dey, A., van Breugel, W., De Vries, W., & Reuland, M. 2002, ApJ, 570, 92
 Dawson, S., Stern, D., Bunker, A. J., Spinrad, H., & Dey, A. 2001, AJ, 122, 598
 Dey, A., Spinrad, H., Stern, D., Graham, J. R., & Chaffee, F. H. 1998, ApJ, 498, L93
 Djorgovski, S. G., Castro, S., Stern, D., & Mahabal, A. A. 2001, ApJ, 560, L5
 Ellis, R., Santos, M. R., Kneib, J. -P., & Kuijken, K. 2001, ApJ, 560, L119
 Fan, X., et al. 2000, AJ, 120, 1167
 Goodrich, R. W., et al. 2001, ApJ, 561, L23
 Heckman, T. M. 1999, After the Dark Ages: When Galaxies were Young (the Universe at $2 < z < 5$). 9th Annual October Astrophysics Conference in Maryland, edited by S. Holt and E. Smith (American Institute of Physics Press), 322
 Hu, E. M., Cowie, L. L., & McMahon, R. G. 1998, ApJ, 502, L99
 Hu, E. M., Cowie, L. L., McMahon, R. G., Capak, R., Iwamuro, F., Kneib, J. -P., Maihara, T., & Motohara, K. 2002, ApJ, 568, L75
 Hu, E. M., & McMahon, R. G. 1996, Nature, 382, 231
 Hu, E. M., McMahon, R. G., & Cowie, L. L. 1999, ApJ, 522, L9
 Kaifu, N. 1998, Proc. SPIE, 3352, 14
 Kashikawa, N., et al. 2000, SPIE, 4008, 104
 Kennicutt, R. C., Jr. 1998, ARA&A, 36, 189
 Kudritzki, R. -P., et al. 2000, ApJ, 536, 19
 Massey, P., Strobel, K., Barnes, J. V., & Anderson, E. 1988, ApJ, 328, 315
 Miyazaki, S., Sekiguchi, M., Imi, K., Okada, N., Nakata, F., & Komiyama, Y. 1998, SPIE, 3355, 363
 Oke, J. B. 1990, AJ, 99, 1621
 Sheinis, A. I., Miller, J. S., Bolte, M., & Sutin, B. 2000, SPIE, 4008, 522
 Spinrad, H., Stern, D., Bunker, A., Dey, A., Lanzetta, K., Yahil, A., Pascarelle, S., & Fernández-Soto, A. 1998, AJ, 116, 2617
 Steidel, C. C., Adelberger, K. L., Shapley, A. E., Pettini, M., Dickinson, M., & Giavalisco, M. 2000, ApJ, 532, 170
 Stern, D., Bunker, A., Spinrad, H., & Dey, A. 2000, ApJ, 537, 73
 Taniguchi, Y., & Shioya, Y. 2000, ApJ, 547, 146
 Weymann, E. J., Stern, D., Bunker, A., Spinrad, H., Chaffee, F. H., Thompson, R. I., & Storrie-Lombardi, L. 1998, ApJ, 505, L95
 Yagi, M., Kashikawa, N., Sekiguchi, M., Doi, M., Yasuda, N., Shimasaku, K. and Okamura, S., 2002, AJ, 123, 66

TABLE 1
JOURNAL OF IMAGING OBSERVATIONS

Band	Obs. Date (UT)	T_{int} (sec) ^a	$m_{\text{lim}}(\text{AB})^b$	$FWHM_{\text{star}}$ (arcsec) ^c
B	2002 February 17	1680	27.1	1.2
R_C	2002 February 15, 16	4800	26.8	1.4
I_C	2002 February 15, 16	3360	26.2	1.2
$NB816$	2002 February 15 - 17	36000	26.6	0.9
z'	2002 February 15, 16	5160	25.4	1.2

^aTotal integration time.

^bThe limiting magnitude (3σ) within a $2''$ aperture.

^cThe full width at half maximum of stellar objects in the final image




FIG. 1.— A finding chart of LAE J1044-0130 in our NB816 image. The field size is $2' \times 2'$. North is up and east is to the left.




FIG. 2.— Thumbnail images of LAE J1044-0130 (upper panel) also displayed as contours (middle panel). The angular size of the circle in each panel corresponds to $8''$. The lower panel shows the spectral energy distribution (in magnitudes). From our optical spectroscopy, we find that a galaxy located at $2''.4$ northeast of LAE J1044-0130 shows two emission lines at $\lambda \sim 6710 - 6720 \text{ \AA}$. Since these lines can be identified as the $[\text{O II}]\lambda 3727$ doublet redshifted to $z = 0.802 \pm 0.002$. This foreground galaxy might enhance the image of LAE J1044-0130 due to gravitational lensing. However, since there is no counter image of the LAE J1044-0130 in our deep NB816 image, the magnification factor might be less than a factor of 2.




FIG. 3.— The optical spectra of LAE J1044-0130 obtained with FOCAS on Subaru. (a) The spectrum shown in the upper panel ($R \sim 400$) was obtained on 2002, March 11. (b) The spectrum shown in lower panel ($R \sim 1000$) was obtained on 2002, March 13.




FIG. 4.— The optical spectrogram (upper panel) and one-dimensional spectrum (middle panel) of LAE J1044-0130 obtained with ESI on Keck II ($R \sim 3400$); note that five-pixel binning was applied to the one-dimensional spectrum. The model profile fit is shown by the thick solid curve (see text). Sky (OH airglow) emission lines are shown in the lower panel.

This figure "f2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207303v2>

This figure "f4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0207303v2>